

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 979 847 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:
16.02.2000 Bulletin 2000/07

(51) Int. Cl.⁷: **C09B 31/16**, C09B 69/02,
C09D 11/02

(21) Application number: 98917653.2

(86) International application number:
PCT/JP98/01877

(22) Date of filing: 23.04.1998

(87) International publication number:
WO 98/47970 (29.10.1998 Gazette 1998/43)

(84) Designated Contracting States:
CH DE FR GB IT LI

(30) Priority: 24.04.1997 JP 12012197

(71) Applicant:
NIPPON KAYAKU KABUSHIKI KAISHA
Tokyo 102-0071 (JP)


(72) Inventors:
• ONO, Hiroaki
Tokyo 174-0051 (JP)
• MIYAZAWA, Takashi
Tokyo 115-0042 (JP)

• TABEL, Toru
Saitama 364-0007 (JP)
• SHIRASAKI, Yasuo
Saitama 330-0826 (JP)
• YAMAGUCHI, Isao
Chiba 270-0142 (JP)
• NAGASAKI, Kazunobu
Saitama 332-0014 (JP)

(74) Representative:
Türk - Gille - Hrabal - Struck
Patentanwälte - European Patent Attorneys,
Brucknerstrasse 20
40593 Düsseldorf (DE)

(54) **WATER-SOLUBLE TRISAZO COMPOUNDS, AQUEOUS INK COMPOSITION AND COLORED ARTICLE**

(57) A black recording liquid having high solubility, being stable even after being stored for long, and capable of forming a printed image with a high optical density and excellent water and light resistances. The recording fluid contains a trisazo compound of general formula (1): wherein D is an 8-hydroxy-1-naphthyl group substituted with one or two sulfo groups; A is a 1,4-phenylene group or 1,4-naphthylene group which may be substituted with amino, hydroxy, alkyl, alkoxy, acylamino or ureido; and B is a phenyl group which may be substituted with hydroxy, amino or sulfo, or a salt thereof and is used for the recording.



EP 0 979 847 A1

Description

Filed of the Invention

5 [0001] The present invention relates to new water-soluble trisazo compounds and colored articles therefrom.

Prior Arts

10 [0002] Among various color recording methods, one typical method is the recording method by means of ink jet printers wherein ink droplets are produced and deposit on various materials to be recorded such as papers, films and fabrics so that recordings are effected. This method has advantages that it is silent since materials to be recorded are not contacted with a recording head and that miniaturization, speeding up and color printing can be easily achieved. Thus, it has recently come into wide use and it is expected to greatly spread in future.

15 [0003] As inks for pens, felt-tip pens and the like as well as inks for ink jet printers, aqueous inks comprising water-soluble dyes dissolved in aqueous media are used. Generally, the aqueous inks comprise water-soluble organic solvents in order to prevent the inks from clogging in nibs and ink delivery nozzles. These inks should meet the requirements that they can form images having a high optical density, that they do not clog nibs and ink delivery nozzles, that they are easily dried on materials to be recorded, that they hardly blot and that they have excellent storage stability. In addition, images therefrom should have sufficient light and water resistances.

20 [0004] Various inks having different hues are prepared from various dyes. Among the inks, black inks are important since they are used in mono-color and full-color images. As dyes for black inks, many dyes have been proposed as described in prior patent applications, for example, JP-55144067A(1980), JP-57207660A(1982), JP-58147470(1983), JP-59093766A(1984), JP-62190269A(1987), JP-62246975A(1987), JP-63022867A(1988), JP-63033484A(1988), JP-01093389A(1989), JP-02140270A(1990), JP-03167270A(1991), JP-03200852A(1991), JP-04359065A(1992), JP-06172668A(1994), JP-06248212A(1994), JP-07026160A(1995), JP-07268256A(1995) and the like. However, the prior
25 dyes cannot satisfy the commercial requirements as described above.

Background of the Invention


30 [0005] An object of the present invention is to provide a black recording fluid having a high solubility, stable even after a long storage, capable of forming a printed image with a high optical density and excellent in water and light resistances.

Summary of the Invention

35 [0006] The present inventors have earnestly investigated in order to solve the aforementioned problem, achieving the present invention.

[0007] Accordingly, the present invention relates to:

40 (1) a trisazo compound represented by the following general formula (1):





50 wherein

D is an 8-hydroxy-1-naphthyl group substituted with one or two sulfo groups;


A is a group of the general formula (2) or (3):

55




wherein R^1 and R^2 are each independently hydrogen, amino, hydroxyl, carboxyl, sulfo, phosphono, (C1-C4)alkoxycarbonyl, (C1-C4)alkyl optionally substituted with hydroxy or (C1-C4)alkoxy, (C1-C4)alkoxy optionally substituted with hydroxy or (C1-C4)alkoxy, (C2-C4)alkanoylamino optionally substituted with hydroxy or (C1-C4)alkoxy or ureido, R^3 is hydrogen, (C1-C4)alkyl or (C1-C4)alkoxy, and R^4 is hydrogen, sulfo or phosphono;

B is a group represented by the general formula (4):



wherein R^5 , R^6 and R^7 are each independently hydrogen, hydroxyl, amino, carboxyl, (C1-C4)alkoxy, (C1-C4)alkoxycarbonyl, (C1-C4)alkyl, mono- or bis-carboxy-(C1-C3)alkylamino, carboxy-(C1-C3)alkoxy, (C1-C4)alkylamino optionally substituted with hydroxy or (C1-C4)alkoxy, (C1-C4)alkanoylamino, sulfo, halogen or ureido; and
 m is 0 or 1,
 or its salt;

(2) a salt of a trisazo compound as defined in (1) wherein the salt is a lithium salt, a sodium salt, or an ammonium salt represented by the general formula (5):



wherein R^8 to R^{11} are each independently hydrogen, (C1-C4)alkyl, hydroxy-(C1-C4)alkyl or hydroxyethoxy-(C1-C4)alkyl group;

(3) a trisazo compound or its salt as defined in (1) or (2) wherein D is

8-hydroxy-3,6-disulfo-1-naphthyl,
 8-hydroxy-4,6-disulfo-1-naphthyl,
 8-hydroxy-2,4-disulfo-1-naphthyl,
 8-hydroxy-3,5-disulfo-1-naphthyl,
 8-hydroxy-4,7-disulfo-1-naphthyl,
 8-hydroxy-5,7-disulfo-1-naphthyl,
 8-hydroxy-4-sulfo-1-naphthyl,
 8-hydroxy-5-sulfo-1-naphthyl, or
 8-hydroxy-6-sulfo-1-naphthyl;

(4) a trisazo compound or its salt as defined in any one of (1) to (3) wherein A is the group of the general formula (2) wherein R¹ and R² are each independently hydrogen, methyl, ethyl, methoxy, ethoxy, methoxyethoxy, amino, hydroxyl, carboxyl, sulfo, acetylamino, n-propionylamino or ureido, or the general formula (3) wherein R³ is hydrogen, methyl or methoxy and R⁴ is hydrogen or sulfo;

(5) a trisazo compound or its salt as defined in any one of (1) to (4) wherein A is the group of the general formula (2) wherein R¹ is methyl, methoxy or methoxyethoxy and R² is hydrogen, methyl, methoxy, amino, hydroxyl, acetylamino or ureido;

(6) a trisazo compound or its salt as defined in any one of (1) to (4) wherein A is the group of the general formula (3) wherein R³ is hydrogen or methoxy and R⁴ is hydrogen or sulfo;

(7) a trisazo compound or its salt as defined in any one of (1) to (6) wherein B is the group of the general formula (4) wherein R⁵ is hydrogen, methyl, methoxy, chlorine, carboxyl or sulfo, and R⁶ and R⁷ are each independently hydroxyl, amino, carboxymethylamino, carboxymethoxy, acetylamino or ureido;

(8) a trisazo compound or its salt as defined in any one of (1) to (7) wherein m is zero;

(9) an aqueous ink composition containing a trisazo compound or its salt as defined in any one of (1) to (8);

(10) an article colored with a trisazo compound as defined in any one of (1) to (8) or an aqueous ink composition as defined in (9); and

(11) a colored article as defined in (10) wherein coloring is effected by means of a printer.

Detailed Description of the Invention

[0008] The present invention will be fully explained below.

[0009] In the compound represented by the aforementioned formula (1), D is an 8-hydroxy-1-naphthyl group substituted with one or two sulfo groups. Preferably, it includes 8-hydroxy-3,6-disulfo-1-naphthyl, 8-hydroxy-4,6-disulfo-1-naphthyl, 8-hydroxy-2,4-disulfo-1-naphthyl, 8-hydroxy-3,5-disulfo-1-naphthyl, 8-hydroxy-4,7-disulfo-1-naphthyl, 8-hydroxy-5,7-disulfo-1-naphthyl, 8-hydroxy-4-sulfo-1-naphthyl, 8-hydroxy-5-sulfo-1-naphthyl, 8-hydroxy-6-sulfo-1-naphthyl and the like. 8-Hydroxy-3,6-disulfo-1-naphthyl, 8-hydroxy-4,6-disulfo-1-naphthyl, 8-hydroxy-2,4-disulfo-1-naphthyl, 8-hydroxy-4-sulfo-1-naphthyl or the like is more preferable.

[0010] In A and B in the aforementioned formula (1), examples of (C1-C4)alkyl include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl or the like, preferably methyl or ethyl. Examples of (C1-C4) alkyl group substituted with hydroxy or (C1-C4)alkoxy include 2-hydroxyethyl, 2-hydroxypropyl, 1-hydroxy-1-methylethyl, methoxyethyl, ethoxyethyl or the like, preferably 2-hydroxyethyl or methoxyethyl. Examples of (C1-C4)alkoxy group optionally substituted with hydroxy or (C1-C4)alkoxy include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, sec-butoxy, tert-butoxy, 2-hydroxyethoxy, 2- or 3-hydroxypropoxy, methoxyethoxy, ethoxyethoxy, n-propoxyethoxy, isopropoxyethoxy, n-butoxyethoxy, methoxypropoxy, ethoxypropoxy, n-propoxypropoxy, isopropoxybutoxy, n-propoxybutoxy, 2-hydroxyethoxyethoxy or the like, preferably methoxy, ethoxy, methoxyethoxy or ethoxyethoxy. Methoxy or methoxyethoxy is particularly preferable. Examples of (C1-C4)alkylamino group optionally substituted with hydroxy or (C1-C4)alkoxy include 2-hydroxyethylamino, 2-hydroxyethylmethylamino, N,N-di(2-hydroxyethyl)amino, 2-hydroxypropylamino, 3-hydroxypropylamino, methoxyethylamino, ethoxyethylamino or the like, preferably 2-hydroxyethylamino, methoxyethylamino or 3-hydroxypropylamino. (C1-C4)alkanoylamino group optionally substituted with hydroxy or (C1-C4)alkoxy include acetylamino, n-propionylamino, isopropionylamino, hydroxyacetylamino, 2- or 3-hydroxy-n-propionylamino, 2- or 3-methoxy-n-propionylamino, butyrylamino or the like, among which acetylamino is particularly preferable. Examples of mono- or bis-carboxy-(C1-C3)alkylamino group include carboxymethylamino, carboxyethylamino, carboxypropylamino, bis-carboxymethylamino, bis-carboxyethylamino, bis-carboxypropylamino, preferably carboxymethylamino, bis-carboxymethylamino or carboxyethylamino. Examples of carboxy-(C1-C3)alkoxy include carboxymethoxy, carboxyethoxy and carboxypropoxy, particularly preferably carboxymethoxy.

[0011] Preferably, R¹ and R² are each independently hydrogen, methyl, ethyl, methoxy, ethoxy, methoxyethoxy, ethoxyethoxy, amino, hydroxyl, carboxyl, sulfo, phosphono, acetylamino, n-propionylamino and ureido. Hydrogen, methyl, methoxy, methoxyethoxy, amino, hydroxyl, carboxyl, acetylamino or ureido is particularly preferable.

[0012] The general formula (2) is preferably derived from each of the following intermediates:

1-amino-2-methoxy-5-methylbenzene, 1-amino-2,5-dimethylbenzene, 1-amino-2-methoxy-5-acetylaminobenzene, 1-amino-2-methoxyethoxy-5-acetylaminobenzene, 1-amino-2-methoxybenzene, 1-amino-2-methylbenzene, 1-amino-2,5-dimethoxybenzene, 1-amino-2-methyl-5-acetylaminobenzene, 1-amino-2-methoxy-5-ureidobenzene, 1-amino-2-methyl-5-ureidobenzene, 1-amino-3-methoxybenzene, 1-amino-3-methylbenzene, 1-amino-2-(2'-hydroxyethoxy)-5-methylbenzene, 3-aminobenzoic acid, 3-amino-4-methylbenzoic acid, 3-amino-4-methoxybenzoic acid, 2,4-diaminobenzoic acid, 2,4-diaminobenzenesulfonic acid, 2-amino-4-acetylaminobenzenesulfonic acid, 3-aminobenzenephosphonic acid, 4-methoxy-3-aminophenol and 2,4-diaminoanisole.

[0013] R^3 is preferably hydrogen, methyl, ethyl, methoxy or ethoxy. Particularly, hydrogen, methyl or methoxy is preferable.

[0014] R^4 is preferably sulfo.

[0015] It is particularly preferable that the residue A of the general formula (3) is derived from the intermediate 1-aminonaphthalene-2-sulfonic acid, 1-aminonaphthalene-3-sulfonic acid, 1-aminonaphthalene-5-sulfonic acid, 1-aminonaphthalene-6-sulfonic acid, 1-aminonaphthalene-7-sulfonic acid, 1-aminonaphthalene-8-sulfonic acid or 1-amino-2-methoxynaphthalene-6-sulfonic acid.


[0016] A component from which the naphthalene residue in the general formula (1) is derived is 1-hydroxy-3-sulfo-7-aminonaphthalene (γ acid) when m is 0 and 1-hydroxy-3,6-disulfo-7-aminonaphthalene (2R acid) when m is 1.

[0017] R^5 is hydrogen, methyl, methoxy, chlorine, carboxyl, sulfo or the like. Methyl, ethyl, methoxy, ethoxy or sulfo is preferable. Particularly, hydrogen or sulfo is preferable.

[0018] R^6 and R^7 are each independently hydroxyl, amino, carboxymethylamino, carboxypropylamino, bis-carboxymethylamino, acetylaminomino, ureido or the like. Hydroxyl, amino, acetylaminomino or carboxymethylamino are particularly preferable.

[0019] Examples of intermediate from which the residue B of the general formula (4) is derived include 1,3-phenylenediamine, resorcin, 3-aminophenol, 2-amino-4-acetylaminobenzenesulfonic acid, 2,4-diaminobenzenesulfonic acid, 2,4-diaminobenzoic acid, 2,4-diaminotoluene, 2,4-diaminoanisole, 2,4-diaminochlorobenzene, 1,3-bis-carboxymethylaminobenzene, 1-amino-3-acetylaminobenzene, 1-amino-2-methoxy-5-acetylaminobenzene, 1-amino-2-methyl-5-acetylaminobenzene, 1-amino-2-methoxy-5-ureidobenzene, 2,4-dihydroxybenzoic acid, 1-amino-2-methyl-5-ureidobenzene or the like. 1,3-Phenylenediamine, resorcin, 3-aminophenol or 2,4-diaminobenzenesulfonic acid is preferable. Their mixture may be used.

[0020] Salts are those of inorganic or organic cations. Suitable inorganic salts are alkali metal or alkaline earth metal salts. Preferable inorganic salt are lithium or sodium salt. And, suitable organic salts are ammonium salts represented by the general formula (5).



[0021] In the above general formula (5), R^8 to R^{11} are each independently hydrogen, (C1-C4)alkyl, hydroxy-(C1-C4)alkyl or hydroxyethoxy-(C1-C4)alkyl. Examples of (C1-C4) alkyl include methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, sec-butyl, tert-butyl or the like. Examples of hydroxy-(C1-C4)alkyl include hydroxymethyl, hydroxyethyl, 3-hydroxypropyl, 2-hydroxypropyl, 4-hydroxybutyl, 3-hydroxybutyl, 2-hydroxybutyl or the like. Examples of hydroxyethoxy-(C1-C4)alkyl include hydroxyethoxymethyl, 2-hydroxyethoxyethyl, 3-hydroxyethoxypropyl, 2-hydroxyethoxypropyl, 4-hydroxyethoxybutyl, 3-hydroxyethoxybutyl, 2-hydroxyethoxybutyl or the like.

[0022] Examples of the compound of the general formula (5) are shown in Table 1.


Table 1

	R^8	R^9	R^{10}	R^{11}
5-1	H	-C ₂ H ₄ OH	-C ₂ H ₄ OH	-C ₂ H ₄ OH
5-2	CH ₃	-C ₂ H ₄ OH	-C ₂ H ₄ OH	-C ₂ H ₄ OH

Table 1 (continued)

	R ⁸	R ⁹	R ¹⁰	R ¹¹
5-3	H	-CH ₂ CH(OH)CH ₃	-CH ₂ CH(OH)CH ₃	-CH ₂ CH(OH)CH ₃
5-4	CH ₃	-CH ₂ CH(OH)CH ₃	-CH ₂ CH(OH)CH ₃	-CH ₂ CH(OH)CH ₃
5-5	H	-C ₂ H ₄ OH	H	-C ₂ H ₄ OH
5-6	CH ₃	-C ₂ H ₄ OH	H	-C ₂ H ₄ OH
5-7	H	-CH ₂ CH(OH)CH ₃	H	-CH ₂ CH(OH)CH ₃
5-8	CH ₃	-CH ₂ CH(OH)CH ₃	H	-CH ₂ CH(OH)CH ₃
5-9	CH ₃	-C ₂ H ₄ OH	CH ₃	-C ₂ H ₄ OH
5-10	CH ₃	-CH ₂ CH(OH)CH ₃	CH ₃	-CH ₂ CH(OH)CH ₃

[0023] Examples of the trisazo compound represented by the general formula (1) are illustrated in the general formula (1-1) (Table 2) to the general formula (1-5) (Table 27).



general formula (1-1)
Compound No.

		A	B
5			
	0001		
	0002	"	
10			
	0003	"	
15	0004	"	
20	0005	"	
	0006	"	
25			
	0007	"	
	0008	"	
30			
	0009	"	
35	0010		
	0011	"	
40			
	0012	"	
45	0013	"	
	0014	"	
50			
	0015	"	

general formula (1-1)
Compound No.

5		A	B
	0016		
10	0017	"	
15	0018	"	
20	0019		
	0020	"	
	0021	"	
25	0022	"	
30	0023		
	0024	"	
35	0025	"	
40	0026	"	
45	0027	"	
	0028	"	
50	0029	"	
55			





















general formula (1-1)
Compound No.

		A	B
5	0030		
	0031	"	
10	0032	"	
	0033	"	
15	0034		
20	0035	"	
	0036	"	
25	0037	"	
	0038		
30	0039	"	
	0040	"	
35	0041	"	
40	0042		
	0043	"	
45	0044	"	

50

55


general formula (1-1)
Compound No.

	A	B
0045		
0046	"	
0047	"	
0048	"	
0049	"	
0050	"	
0051		
0052		
0053	"	
0054	"	
0055	"	
0056		
0057	"	
0058		
0059	"	


general formula (1-1)
Compound No.

5

0060




B



10


0061

"



0062


"



15



0063

"




20

0064



0065


"



25



0066

"




30

0067



0068


"



35



0069

"




40

0070





0071

"




45

0072



0073

"




50


55

general formula (1-1)
Compound No.


5



10




15






20




25





30




35



40




45




















50

55






















general formula (1-2)
Compound No.

		A	B
5	0086		
	0087	"	
10	0088	"	
15	0089	"	
20	0090	"	
	0091	"	
25	0092	"	
30	0093	"	
	0094	"	
35	0095		
	0096	"	
40	0097	"	
45	0098	"	
	0099	"	
50	0100	"	

general formula (1-2)
Compound No.

		A	B
5	0101		
	0102	"	
10	0103	"	
	0104		
15	0105	"	
20	0106	"	
	0107	"	
25	0108		
30	0109	"	
35	0110	"	
	0111	"	
40	0112	"	
45	0113	"	
50	0114	"	





















general formula (1-2)
Compound No.

5		A	B
	0115		
10	0116	"	
	0117	"	
15	0118	"	
	0119		
20	0120	"	
25	0121	"	
	0122	"	
30	0123		
	0124	"	
35	0125	"	
	0126	"	
40	0127		
45	0128	"	
	0129	"	

50

55

general formula (1-2)
Compound No.

		A	B
5	0130		
	0131	"	
10	0132	"	
	0133	"	
15	0134	"	
20	0135	"	
	0136		
25	0137		
30	0138	"	
	0139	"	
35	0140		
	0141	"	
40	0142	"	
45	0143		
50	0144	"	



general formula (1-2)

Compound No.

A


B

0145




0146

"




0147

"





0148

"




0149




0150

"





0151

"




0152




0153

"





0154

"




0155





0156

"




0157



0158

"





general formula (1-2)
Compound No.

A


B

0159





0160

"




0161





0162

"




0163





0164



"





0165





0166




0167





0168




0169



0170





10

general formula (1-3)
Compound No.

Compound No.	A	B
0171		
0172	"	
0173	"	
0174	"	
0175		
0176	"	
0177	"	
0178	"	
0179	"	
0180	"	
0181		
0182	"	
0183	"	



general formula (1-3)

Compound No.

A


B

0184




0185

"





0186

"




0187




0188

"





0189

"




0190





0191

"




0192




0193

"




0194

"



0195

"





general formula (1-3)
Compound No.

A


B

0196




0197

"




0198

"





0199

"




0200





0201

"




0202





0203

"




0204




0205

"





0206

"




0207





0208

"




0209



0210

"





general formula (1-3)

Compound No.

A


B

0211





0212

"




0213





0214

"




0215





0216



"





0217






0218



0219



0220



(1-4)



general formula (1-4)
Compound No.

A

B


5

0221



0222


"



10


0223

"



0224


"



15

0225


"



20

0226


"



25


0227

"



0228


"



30



0229

"




35

0230



0231


"



40

0232


"



45


0233

"



0234


"



50

0235

"




55


general formula (1-4)
Compound No.

5

0236




B



10


0237

"





0238

"




15

0239



0240


"



20

0241


"



25



0242

"




30

0243



0244


"



35

0245


"



40


0246

"



0247


"



45

0248


















"






















50

55




















general formula (1-4)
Compound No.

5		A	B
	0249		
10	0250	"	
	0251	"	
15	0252	"	
20	0253		
	0254	"	
25	0255	"	
30	0256	"	
	0257		
35	0258	"	
40	0259	"	
45	0260	"	
	0261	"	
50	0262	"	
55			

general formula (1-4)
Compound No.

		A	B
5	0263		
10	0264	"	
	0265	"	
15	0266	"	
	0267	"	
20	0268	"	
25	0269		
	0270		
30	0271	"	
	0272	"	
35	0273		
40	0274	"	
	0275		
45	0276	"	

general formula (1-4)
Compound No.

		A	B
5	0277		
10	0278	"	
	0279	"	
15	0280	"	
20	0281		
	0282	"	
25	0283	"	
30	0284		
	0285	"	
35	0286	"	
40	0287		
	0288	"	
45	0289		
50	0290	"	
55			



general formula (1-4)
Compound No.

A

B



5

0291




10

0292





0293

"




15

0294





0295

"





20

0296





25

0297





30

0298



35


0299





40

0300

"




0301



45




















50



(1-5)

55


general formula (1-5)
Compound No.

		A	B
5	0302		
10	0303	"	
	0304	"	
15	0305	"	
20	0306	"	
	0307		
25	0308	"	
	0309	"	
30	0310	"	
35	0311	"	
40	0312	"	 + 
	0313	"	
45	0314		
	0315	"	
50	0316	"	


general formula (1-5)
Compound No.

5

0317




B



10


0318

"



0319


"



15



0320

"



20


0321



25


0322

"





0323

"



30


0324





35

0325

"




0326



40


0327

"



0328


"



45

0329


















"



50

55

general formula (1-5)
Compound No.


5		A	B
	0330		
10	0331	"	
	0332	"	
15	0333	"	
20	0334	"	
	0335		
25	0336	"	
30	0337		
	0338	"	
35	0339	"	
40	0340		
	0341	"	
45	0342	"	
50			
55			

general formula (1-5)
Compound No.

	A	B
0343		
0344	"	
0345		
0346	"	
0347		
0348		
0349	"	
0350		
0351		
0352		
0353		
0354	"	


[0024] The compound represented by the general formula (1) can be synthesized, for example, according to the following method. That is, the compound of the general formula (1) can be synthesized by:

-- diazotizing a compound of the general formula (6):




10
wherein Y is hydrogen, halogen or (C1-C4) alkyl and A is as defined above,
in an aqueous medium;

15
-- coupling with a compound of the general formula (7):




25
wherein m is 0 or 1;

-- hydrolyzing the resultant disazo compound of the general formula (8):




35
wherein A, Y and m are as defined above, to a compound of the general formula (9):



45
wherein A and m are as defined above;

-- diazotizing the compound of the general formula (9); and

50
-- coupling with a compound of the general formula (4):



wherein R^5 , R^6 and R^7 are as defined above. Alternatively, the compound of the general formula (8) may be diazotized again, coupled with the compound of the general formula (4') and then hydrolyzed to obtain the compound of the general formula (1).

[0025] Each of the compounds of the general formulae (6), (7) and (4') can be prepared according to the known method per se.

[0026] Diazotization of the compound of the general formula (6) is effected according to the known method per se, for example, using nitrites such as an alkali metal nitrite (e.g. sodium nitrite) at -5 to 30°C , preferably 0 to 10°C in an aqueous inorganic acid medium.

[0027] Coupling of the diazo compound of the general formula (6) with the compound of the general formula (7) is effected under the known condition per se. It is conveniently effected in an aqueous or aqueous organic medium at -5 to 30°C , preferably 0 to 10°C under the weak acidic or alkaline pH. Preferably, it is effected under the neutral or alkaline pH, e.g. 7 to 11 . The pH is adjusted by the addition of any base. The base to be used includes alkali metal hydroxides such as lithium hydroxide and sodium hydroxide; alkali metal carbonates such as lithium carbonate, sodium carbonate and potassium carbonate; alkali metal acetates such as sodium acetate; ammonia; or organic amines. The compounds of the general formulae (6) and (7) are used in approximately stoichiometric amounts in this reaction.

[0028] The preparation of the compound of the general formula (9) by hydrolyzing the compound of the general formula (8) is effected according to the known method per se. The heating in an aqueous alkaline medium is convenient. For example, it is effected by adding sodium or potassium hydroxide to a reaction solution containing the compound of the general formula (8) until the pH is 11 or more and then heating to the temperature of 20 to 150°C , preferably 30 to 100°C . Preferably, the pH of the reaction solution is kept to be 10 to 12 .

[0029] Diazotization of the compound of the general formula (9) resulting from hydrolyzing is effected under the same condition as that described in the diazotization of the compound of the general formula (6).

[0030] Coupling of the diazotized product of the compound of the general formula (9) with the compound of the general formula (4') is effected under the known condition per se. It is effected in an aqueous or aqueous organic medium at the temperature of 0 to 30°C , preferably 10 to 20°C and the pH of 4 to 10 , preferably 6 to 8 .

[0031] After the coupling reaction, the compound represented by the general formula (1) of the present invention can be isolated in the form of free acid by adding any mineral acid. The thus-isolated free acid may be washed with water or an acidified water to remove inorganic salts. The acid type dye having a low salt content can be converted to a solution containing its corresponding salt by neutralizing with hydroxide and/or carbonate (e.g. LiOH , Li_2CO_3 , NaOH , Na_2CO_3 or K_2CO_3) or amine in an aqueous medium. Preferable examples of the amine include ammonia, ethanolamine, diethanolamine, triethanolamine, N-methyl-N-ethanolamine, N-methyl-N,N-diethanolamine, 2-(2-hydroxyethoxy)-ethylamine, di-2-(2-hydroxyethoxy)-ethylamine, tri-2-(2-hydroxyethoxy)-ethylamine or the like.

[0032] The trisazo compound represented by the general formula (1) is suitable for dyeing papers, natural fibers such as celluloses, wools and leathers and blended yarn fabrics of natural and synthetic fibers. Further, it is suitable for the preparation of writing inks and recording liquid.

[0033] The reaction liquid containing the trisazo compound represented by the general formula (1) of the present invention can be used as it is in the preparation of recording ink compositions. Alternatively, the above compound is dried, for example, spray-dried and then processed into an ink composition.

[0034] As a finished product, the recording ink composition contains the trisazo compound represented by the general formula (1) of the present invention in an amount of, for example, about 0.1 to 20% by weight, preferably about 1 to 10% by weight, more preferably about 2 to 8% by weight.

[0035] The ink composition of the present invention may comprises a water-soluble organic solvent in an amount of, for example, 0 to 30% by weight and an ink formulating agent in an amount of, for example, 0 to 5% by weight.

[0036] The aqueous ink composition of the present invention comprises the trisazo compound represented by the general formula (1) dissolved in water or an aqueous solvent (water containing any water-soluble organic solvent or any water-compatible organic solvent). When this aqueous ink composition is used as an ink for ink jet printers, the trisazo compounds having less amounts (for example, about 1% by weight or less) of inorganic substances such as chlorides

and sulfates of metallic cations are preferably used. Such trisazo compounds having less amounts of the inorganic substances may be prepared by subjecting to the desalting treatment according to any conventional method such as the method using a reverse osmotic membrane.

[0037] Examples of water-soluble organic solvents include (C1-C4)alkanol such as methanol, ethanol, propanol, isopropanol, butanol, isobutanol, sec-butanol and tert-butanol; carboxamides such as N,N-dimethylformamide and N,N-dimethylacetamide; lactams such as ϵ -caprolactam and N-methylpyrrolidin-2-one; cyclic ureas such as 1,3-dimethylimidazolin-2-one and 1,3-dimethylhexahydropyrimid-2-one; ketones and ketoalcohols such as acetone, methyl ethyl ketone and 2-methyl-2-hydroxypentan-4-one; cyclic ethers such as tetrahydrofuran and dioxane; mono-, oligo- or polyalkylene glycols or thioglycols having (C2-C6)alkylene units such as ethylene glycol, 1,2- or 1,3-propylene glycol, 1,2- or 1,4-butylene glycol, 1,6-hexylene glycol, diethylene glycol, triethylene glycol, dipropylene glycol, thiodiglycol, polyethylene glycol and polypropylene glycol; polyol (triol) such as glycerol and hexane-1,2,6-triol; (C1-C4)alkyl ethers of polyhydric alcohols such as ethylene glycol monomethyl ether or monoethyl ether, diethylene glycol monomethyl ether or monoethyl ether and triethylene glycol monomethyl ether or monoethyl ether; γ -butyrolactone; disulfoxide and the like.

[0038] Examples of ink formulating agents include antimicrobial and antiseptic agents, pH adjusters, chelating agents, antirusting agents, water-soluble UV absorbing agents, water-soluble polymeric compounds, dye solubilizing agents, surfactants and the like.

[0039] Examples of antimicrobial and antiseptic agents include anhydrous sodium acetate, sodium sorbate, sodium 2-pyridinethiol-1-oxide, sodium benzoate, sodium pentachlorophenol and the like.

[0040] As pH adjusters, any substances can be used as long as they can adjust pH of an ink within the suitable range, for example, in the range from 8.0 to 11.0 without adversely affecting the ink to be formulated. Examples thereof include alkanolamines such as diethanolamine and triethanolamine; alkali metal hydroxides such as lithium hydroxide, sodium hydroxide and potassium hydroxide; ammonium hydroxide; alkali metal carbonates such as lithium carbonate, sodium carbonate and potassium carbonate; and the like.

[0041] Examples of chelating agents include sodium ethylenediamine tetraacetate, sodium nitrilotriacetate, sodium hydroxyethylethylene diamine triacetate, sodium diethylenetriaminepentacetate, sodium uracil diacetate and the like.

[0042] Examples of antirusting agents include acidic sulfite, sodium thiosulfate, ammonium thioglycolate, diisopropylammonium nitrite, pentaerythritol tetranitrate, dicyclohexylammonium nitrite and the like.

[0043] Examples of water-soluble polymeric compounds include polyvinyl alcohol, cellulose derivatives, polyamine, polyimine and the like.

[0044] Examples of water-soluble UV absorbing agents include sulfonated benzophenone, sulfonated benzotriazole and the like.

[0045] Examples of dye solubilizing agents include ϵ -caprolactam, ethylene carbonate, urea and the like.

[0046] Examples of surfactants include any known surfactants of anionic, cationic and nonionic types.

[0047] The colored article of the present invention means any article colored with the compound of the present invention or the aqueous ink composition containing said compound. Examples of the articles to be colored include, but not to be limited to, papers, fibers and clothes (cellulose, nylon, wool and the like), leathers, substrates for color filter and the like. Any coloring methods which can be mentioned are the printing method such as the dip dyeing, the textile printing, or the screen printing and the method by means of ink jet printers. Among these coloring methods, printing with an ink jet printers is preferred.

[0048] Suitably, the aqueous ink composition containing the compound represented by the general formula (1) and optionally any other compound (dye) is useful for printing, copying, marking, writing, drawing, stamping and recording, particularly ink jet printing. Thereby, black prints of high quality having a remarkably full depth of shade and good resistances properties to water, light and abrasion can be obtained. The trisazo compound represented by the general formula (1) of the present invention exhibits improved color fastness to light, especially when applied onto papers intended for ink jet printing.


[0049] The aqueous ink composition of the present invention does not produce precipitates during its storage. In case the aqueous ink composition of the present invention is used in ink jet printing, it does not cause the clogging of a jet nozzle. And, when continuously used in the ink jet printer with certain recirculation for considerably long time or intermittently used in the ink jet printer, the aqueous ink composition of the present invention shows no change in physical properties.

Examples


[0050] The following examples will more fully illustrate the embodiments of the invention. All parts and percentages referred to herein are by weight unless otherwise indicated.

Example 1

[0051] A solution prepared by dissolving 62.1 parts of the compound of formula (10) in 500 parts of water under a weak alkaline condition and then adding 10.3 parts of sodium nitrite was added dropwise to 400 parts of 4.5% hydrochloric acid at 5 to 10°C to diazotize.



[0052] The suspension of the resultant diazo of the compound (10) was then added dropwise to an aqueous solution of 22.9 parts of 7-amino-1-hydroxynaphthalene-3-sulfonic acid dissolved in 500 parts of water at 0 to 5°C over about 30 minutes. During this addition, the pH of the coupling was kept to be 8.5 to 9.5 by adding an aqueous sodium carbonate solution. After the addition, the reaction mixture was stirred at 5 to 10°C for 3 hours and at 15 to 20°C for 3 hours while keeping the pH of 8.5 to 9.0 to complete the coupling reaction. After an aqueous sodium hydroxide solution was added until the pH was 12 to 13, the reaction mixture was stirred at 80 to 95°C and at the pH 10 to 11 for about 2 hours. Then, the solution was allowed to be cooled to room temperature and neutralized with concentrated hydrochloric acid until the pH was 6 to 7, to which sodium chloride was added to salt out. Precipitates were isolated by filtration and dried to obtain 62.4 parts of the compound of formula (11).



[0053] 7.2 Parts of the compound of formula (11) thus obtained were dissolved in 150 parts of hot water, to which 0.83 part of sodium nitrite was added. The solution was added dropwise to 50 parts of 4.5 % hydrochloric acid at 10 to 15°C to diazotize. After 2.0 parts of 2,4-diaminobenzene-sulfonic acid was added to the suspension of the resultant diazo compound (11) at the same temperature, the reaction suspension was neutralized with an aqueous sodium carbonate solution over about 1 hour until the pH was 6.5. After the neutralization, it was stirred at 10 to 15°C for 3 hours and at 15 to 20°C for 3 hour while keeping 7 to 8 of pH to complete the coupling reaction. After concentrated hydrochloric acid was added until the pH was 1.0, sodium chloride was added to the reaction mixture to salt out. The thus-precipitated dye was isolated by filtration and dried to obtain 8.2 parts of the black trisazo compound (Compound No. 0005). Absorption spectrum of this compound in water was $\lambda_{\text{max}} = 591 \text{ nm}$.

Example 2

[0054] The compound of formula (11) obtained in Example 1 was diazotized under the same condition as that described in Example 1. After 1.2 parts of 3-aminophenol was added to the suspension of the resultant diazonium salt, the reaction mixture was neutralized with an aqueous sodium carbonate solution over about 1 hour until the pH was 6.5. After the neutralization, it was stirred at 10 to 15°C for 3 hours and at 15 to 20°C for 3 hours while keeping the pH of 7 to 8 to complete the coupling reaction, to which sodium chloride was added to salt out. The thus-precipitated dye was isolated by filtration and dried to obtain 7.9 parts of the black trisazo compound (Compound No. 0002). Absorption spectrum of this compound in water was $\lambda_{\text{max}} = 598 \text{ nm}$.

Example 3

[0055] Coupling was effected under the same condition as that described in Example 2 except that 3-aminophenol was replaced with 1.1 parts of m-phenylenediamine to obtain 8.1 parts of the black trisazo compound (Compound No. 0001). Absorption spectrum of this compound in water was $\lambda_{\max} = 594 \text{ nm}$.

Example 4


[0056] Coupling was effected under the same condition as that described in Example 2 except that 3-aminophenol was replaced with 2.3 parts of N,N'-dicarboxymethyl-m-phenylenediamine to obtain 9.0 parts of the black trisazo compound (Compound No. 0007). Absorption spectrum of this compound in water was $\lambda_{\max} = 582 \text{ nm}$.

Example 5


[0057] Coupling was effected under the same condition as that described in Example 2 except that 3-aminophenol was replaced with 1.2 parts of resorcin to obtain 8.2 parts of the black trisazo compound (Compound No. 0003). Absorption spectrum of this compound in water was $\lambda_{\max} = 597 \text{ nm}$.

Example 6

[0058] A solution prepared by dissolving 75.1 parts of the compound of formula (12) in 500 parts of water under a weak alkaline condition and then adding 10.3 parts of sodium nitrite was added dropwise to 400 parts of 4.5% hydrochloric acid at 5 to 10°C to diazotize.



[0059] The suspension of the resultant diazo compound (12) was added dropwise to an aqueous solution of 22.9 parts of 7-amino-1-hydroxynaphthalene-3-sulfonic acid dissolved in 500 parts of water at 0 to 5°C over about 30 minutes. During this addition, the pH of the coupling was kept to be 8.5 to 9.5 by adding an aqueous sodium carbonate solution. After the addition, the reaction mixture was stirred at 5 to 10°C for 3 hours and at 15 to 20°C for 3 hours while keeping the pH of 8.5 to 9.0 to complete the coupling reaction. After a sodium hydroxide solution was added until the pH was 12 to 13, the reaction mixture was stirred at 80 to 85°C for about 2 hours. Then, the solution was allowed to be cooled to room temperature and neutralized with concentrated hydrochloric acid until the pH was 6 to 7, to which sodium chloride was added to salt out. Precipitates were isolated by filtration and dried to obtain 68.2 parts of the compound of formula (13).




[0060] 8.0 Parts of the compound of formula (13) thus prepared were dissolved in 100 parts of hot water, to which 0.83 part of sodium nitrite was added. The solution was added dropwise to 50 parts of 4.5 % hydrochloric acid at 15 to 20°C to diazotize. After 2.1 parts of 2,4-diaminobenzene-sulfonic acid was added to the suspension of the resultant diazo compound (13) at the same temperature, the reaction suspension was neutralized with an aqueous sodium carbonate solution over about 1 hour until the pH was 6.5. After the neutralization, it was stirred at 10 to 15°C for 3 hours and at 15 to 20°C for 3 hour while keeping the pH of 7 to 8 to complete the coupling reaction. After concentrated hydrochloric acid was added until pH was strongly acidic, the thus-precipitated dye was isolated by filtration and washed with water to remove inorganic salts. Then, the wet cake of the dye was suspended in 500 ml of water and the resultant dye paste was neutralized with 70 ml of a 5N lithium hydroxide solution. The thus-prepared dye solution was spray-dried to obtain 8.7 parts of the lithium salt of the black trisazo compound (Compound No. 0027). Absorption spectrum of this salt in water was $\lambda_{\max} = 604 \text{ nm}$.

Example 7


[0061] Coupling was effected under the same condition as that described in Example 6 except that 2,4-diaminobenzenesulfonic acid was replaced with 1.2 parts of resorcin to obtain 8.2 parts of the black trisazo compound (Compound No. 0025). Absorption spectrum of this compound in water was $\lambda_{\max} = 606 \text{ nm}$.

Example 8

[0062] A solution prepared by dissolving 66.4 parts of the compound of formula (14) in 500 parts of water under a weak alkaline condition and then adding 10.3 parts of sodium nitrite was added dropwise to 400 parts of 4.5% hydrochloric acid at 5 to 10°C to diazotize.



[0063] The suspension of the resultant diazo compound of formula (14) was added dropwise to an aqueous solution of 22.9 parts of 7-amino-1-hydroxynaphthalene-3-sulfonic acid dissolved in 500 parts of water at 0 to 5°C over about 30 minutes. During this addition, the pH of the coupling was kept to be 8.5 to 9.5 by adding an aqueous sodium carbonate solution. After the addition, the reaction mixture was stirred at 5 to 10°C for 3 hours and at 15 to 20°C for 3 hours while keeping the pH of 8.5 to 9.0 to complete the coupling reaction. After a sodium hydroxide solution was added until the pH was 11 to 12, the reaction mixture was stirred at 70 to 75°C for about 4 hours. Then, the solution was allowed to be cooled to room temperature and neutralized with concentrated hydrochloric acid until the pH was 6 to 7, to which sodium chloride was added to salt out. Precipitates were isolated by filtration and dried to obtain 69.5 parts of the compound of formula (15).



[0064] 7.9 Parts of the compound of formula (15) thus prepared were dissolved in 100 parts of hot water, to which 0.83 part of sodium nitrite was added. The solution was added dropwise to 50 parts of 4.5 % hydrochloric acid at 10 to 15°C to diazotize. After 2.1 parts of 2,4-diaminobenzene-sulfonic acid was added to the suspension of the resultant diazo of the compound (15) at the same temperature, the reaction suspension was neutralized with a sodium carbonate solution over about 1 hour until the pH was 6.5. After the neutralization, it was stirred at 10 to 15°C for 3 hours and at 15 to 20°C for 3 hour while keeping the pH of 7 to 8 to complete the coupling reaction. Concentrated hydrochloric acid was added until the pH was 1.0, to which sodium chloride was added to salt out. The thus-precipitated dye was isolated by filtration and dried to obtain 8.9 parts of the black trisazo compound (Compound No. 0014). Absorption spectrum of this compound in water was $\lambda_{\max} = 590$ nm.

Example 9

[0065] The black trisazo compound (Compound No. 0014) obtained in Example 8 was heated in 80 parts of an aqueous 10% hydrochloric acid solution at 80 to 90°C for 6 hours to obtain 6.5 parts of the black trisazo compound (Compound No. 0048). Absorption spectrum of this compound in water was $\lambda_{\max} = 620$ nm.

Example 10


[0066] The compound of formula (15) obtained in Example 8 was diazotized under the same condition as that described in Example 8. After adding 1.2 parts of 3-aminophenol to the suspension of the resultant diazonium salt, the reaction suspension was neutralized with a sodium carbonate solution over about 1 hour until the pH was 6.5. After the neutralization, it was stirred at 10 to 15°C for 3 hours and at 15 to 20°C for 3 hours while keeping the pH of 7 to 8 to complete the coupling reaction, to which sodium chloride was added to salt out. The thus-precipitated dye was isolated by filtration and dried to obtain 7.9 parts of the black trisazo compound (Compound No. 0011). Absorption spectrum of this compound in water was $\lambda_{\max} = 596$ nm.

Example 11

[0067] Coupling was effected under the same condition as that described in Example 8 except that 2,4-diaminobenzenesulfonic acid was replaced with 1.2 parts of resorcin to obtain 8.2 parts of the black trisazo compound (Compound No. 0012). Absorption spectrum of this compound in water was $\lambda_{\max} = 602$ nm.


Example 12

[0068] A solution prepared by dissolving 70.8 parts of the compound of formula (16) in 500 parts of water under a weak alkaline condition and then adding 10.3 parts of sodium nitrite was added dropwise to 400 parts of 4.5% hydrochloric acid at 5 to 10°C to diazotize.



[0069] The suspension of the resultant diazo compound (16) was added dropwise to an aqueous solution of 22.9 parts of 7-amino-1-hydroxynaphthalene-3-sulfonic acid dissolved in 500 parts of water at 0 to 5°C over about 30 minutes. During this addition, the pH of the coupling was kept to be 8.5 to 9.5 by adding an aqueous sodium carbonate solution. After the addition, the reaction mixture was stirred at 5 to 10°C for 3 hours and at 15 to 20°C for 3 hours while keeping the pH of 8.5 to 9.0 to complete the coupling reaction. After a sodium hydroxide solution was added until the pH was 11 to 12, the reaction mixture was stirred at 70 to 75°C for about 4 hours. Then, the solution was allowed to be cooled to room temperature and neutralized with concentrated hydrochloric acid until the pH was 6 to 7, to which sodium chloride was added to salt out. Precipitates were isolated by filtration and dried to obtain 72.3 parts of the com-

pound of formula (17).



15 [0070] 8.0 Parts of the compound of formula (17) thus prepared were dissolved in 100 parts of hot water, to which 0.83 part of sodium nitrite was added. The solution was added dropwise to 50 parts of 4.5 % hydrochloric acid at 10 to 15°C to diazotize. After 2.1 parts of 2,4-diaminobenzene-sulfonic acid was added to the suspension of the resultant diazo compound (17) at the same temperature, the reaction suspension was neutralized with a sodium carbonate solution over about 1 hour until the pH was 6.5. After the neutralization, it was stirred at 10 to 15°C for 3 hours and at 15 to 20°C for 3 hour while keeping the pH of 7 to 8 to complete the coupling reaction. After concentrated hydrochloric acid was added until the pH was 1.0, sodium chloride was added to salt out. The thus-precipitated dye was isolated by filtration and dried to obtain 8.9 parts of the black trisazo compound (Compound No. 0022). Absorption spectrum of this compound in water was $\lambda_{\max} = 591 \text{ nm}$.

25 Example 13

[0071] 8.0 Parts of the black trisazo compound (Compound No. 0022) obtained in Example 12 was heated in 80 parts of an aqueous 10% hydrochloric acid solution at 80 to 90°C for 6 hours to obtain 7.0 parts of the black trisazo compound (Compound No. 0051). Absorption spectrum of this compound in water was $\lambda_{\max} = 620 \text{ nm}$.

30 Example 14


[0072] Coupling was effected under the same condition as that described in Example 12 except that 2,4-diaminobenzenesulfonic acid was replaced with 1.2 parts of 3-aminophenol to obtain 8.0 parts of the black trisazo compound (Compound No. 0020). Absorption spectrum of this compound in water was $\lambda_{\max} = 598 \text{ nm}$.

35 Example 15

[0073] Coupling was effected under the same condition as that described in Example 12 except that 3-aminophenol was replaced with 1.1 parts of m-phenylenediamine to obtain 7.9 parts of the black trisazo compound (Compound No. 0019). Absorption spectrum of this compound in water was $\lambda_{\max} = 596 \text{ nm}$.

Example 16

45 [0074] A solution prepared by dissolving 66.4 parts of the compound of formula (14) in 500 parts of water under a weak alkaline condition and then adding 10.3 parts of sodium nitrite was added dropwise to 400 parts of 4.5% hydrochloric acid at 10 to 15°C to diazotize. The suspension of the resultant diazo compound (14) was added dropwise to an aqueous solution of 31.9 parts of 7-amino-1-hydroxynaphthalene-3,6-disulfonic acid dissolved in 500 parts of water at 5 to 10°C over about 1 hour. During this addition, the pH was kept to be 8.5 to 9.5 by adding an aqueous sodium carbonate solution. After the addition, the reaction mixture was stirred at 10 to 15°C for 3 hours and at 20 to 25°C for 3 hours while keeping the pH of 9.0 to 9.5 to complete the coupling reaction. After a sodium hydroxide solution was added until the pH was 11 to 12, the reaction was stirred at 70 to 75°C for about 4 hours. Then, the solution was allowed to be cooled to room temperature and neutralized with concentrated hydrochloric acid until the pH was 4 to 5, to which sodium chloride was added to salt out. Precipitates were isolated by filtration and dried to obtain 73.8 parts of the compound of formula (18).



(18)

[0075] 8.4 Parts of the compound of formula (18) thus prepared were dissolved in 100 parts of hot water, to which 0.83 part of sodium nitrite was added. The solution was then added dropwise to 50 parts of 4.5 % hydrochloric acid at 20 to 25°C to diazotize. After 2.1 parts of 2,4-diaminobenzene-sulfonic acid was added to the suspension of the resultant diazo compound (18) at the same temperature, the reaction suspension was neutralized with a sodium carbonate solution over about 1 hour until the pH was 6.5. After the neutralization, it was stirred at 20 to 25°C for 5 hours while keeping the pH of 7 to 8 to complete the coupling reaction. After concentrated hydrochloric acid until the pH was 4.0, sodium chloride was added to salt out. The thus-precipitated dye was isolated by filtration and dried to obtain 9.2 parts of the black trisazo compound (Compound No. 0234). Absorption spectrum of this compound in water was $\lambda_{\max} = 605$ nm.

Example 17

[0076] The compound of formula (18) obtained in Example 16 was diazotized under the same condition as that described in Example 16 to diazotize. After adding 1.2 parts of 3-aminophenol, the suspension of the resultant diazonium salt was neutralized with a sodium carbonate solution over about 1 hour until the pH was 6.5. After the neutralization, it was stirred at 20 to 25°C for 5 hours while keeping the pH of 7 to 8 to complete the coupling reaction. Then, sodium chloride was added to salt out. The thus-precipitated dye was isolated by filtration and dried to obtain 7.9 parts of the black trisazo compound (Compound No. 0231). Absorption spectrum of this compound in water was $\lambda_{\max} = 606$ nm.

Example 18

[0077] Coupling was effected under the same condition as that described in Example 16 except that 2,4-diaminobenzenesulfonic acid was replaced with 1.2 parts of resorcin to obtain 8.2 parts of the black trisazo compound (Compound No. 0232). Absorption spectrum of this compound in water was $\lambda_{\max} = 608$ nm.

Example 19

[0078] The black trisazo compound (Compound No. 0231) obtained in Example 17 was heated in 80 parts of an aqueous 10 % hydrochloric acid solution at 80 to 90°C for 6 hours to obtain 6.5 parts of the black trisazo compound (Compound No. 0264). Absorption spectrum of this compound in water was $\lambda_{\max} = 624$ nm.

Example 20

(A) Preparation of ink

[0079] A liquid having the following formulation was prepared and passed through a 0.45 μm membrane filter to obtain an aqueous ink composition for ink jet printers.

(formulation)	
	parts
trisazo compound obtained in each of Examples 1 to 19 (used after desalting)	3.0

(continued)

(formulation)	
	parts
5 water	67.0
ethylene glycol	5.0
glycerol	5.0
10 N-methylpyrrolidone	7.0
urea	10.0
IPA	3.0

15 (B) Ink jet printing

[0080] An ink jet recording was conducted on three different papers, i.e. plain papers (Canon printer paper A4 (TLB5A4S)), ink jet papers A (Color BJ Paper LC 101; ex. Canon) and ink jet papers B (coated papers for ink jet printer STX73 A4; ex. Sharp) by means of an ink jet printer (trade name NOVA JET III; ex. ENCAD). The thus-recorded images
20 were tested with respect to the resistances to light and water and a color rendering property.

(C) Test for resistance to light of the recorded images

[0081] The recorded images were irradiated with light for 20 hours by means of a carbon arc fade meter (ex. SUGA Shikenkisha). Light fastness was judged according to the Blue Scale for light resistance standard as defined in JIS L-0841.
25

(D) Test for resistance to water of the recorded images

[0082] Test specimen was immersed in water contained in a beaker with mild stirring for 2 minutes and dried. Amount of the compound was observed by the visual examination before and after the immersion, and dropout of the compound was judged.
30

- ☐ Less amount of the compound was dropped out.
☐ About half of the compound was dropped out.
☐ Large amount of the compound was dropped out.
35

(E) Color rendering property

[0083] Discoloration was judged by the visual examination by comparing the hue change under the tungsten light with that under the standard light source.
40

- ☐ Change in hue was small.
☐ Change in hue was somewhat large.
☐ Change in hue was large.
45

Comparative Examples 1 and 2


[0084] Two types of dyes, one being used as black dye for ink jet printers and other being proposed as black dye for ink jet printers were tested by way of comparative examples.
50

(1) Comparative Example 1

[0085] An ink was prepared from Cl. Food Black 2 in the same manner as that described in the aforementioned (A) and (B) and used for ink jet printing.
55

(2) Comparative Example 2

[0086] An ink was prepared from the dye represented by the formula (19) in the same manner as that described in the aforementioned (A) and (B) and used for ink jet printing.



[0087] The results are shown in Table 28.

Table 28

Compound No.	light fastness (rank)			water fastness			color rendering property
	plain paper	ink jet paper A	ink jet paper B	plain paper	ink jet paper A	ink jet paper B	
0001	5	4-5	5	Δ~○	○	○	○
0002	4-5	4	5	Δ~○	○	○	○
0003	4-5	3-4	4-5	Δ	○	○	○
0005	5	4-5	5	Δ~○	○	○	○
0007	4	3-4	4	○	○	○	○
0011	4-5	4	4	Δ~○	○	○	○
0012	4-5	3-4	4	Δ	○	○	○
0014	5	5	5	Δ	○	○	○
0019	5	5	5	Δ~○	○	○	○
0020	4-5	4	5	Δ~○	○	○	○
0022	5	5	5	Δ~○	○	○	○
0025	4-5	3	4	X~Δ	○	○	○
0027	5	4	5	X~Δ	○	○	○
0048	4-5	4	5	○	○	○	○
0051	4-5	4	5	○	○	○	○
0231	4-5	4	5	Δ	○	○	○
0232	4	4	4	Δ~○	○	○	○
0234	5	5	5	Δ	○	○	○
0264	4-5	4	5	○	○	○	○
Comp.Ex.1	4-5	3	4-5	X	○	X	Δ
Comp.Ex.2	4	2	4	○	○	○	X

[0088] As shown in Table 28, the ink composition containing the compound of the present invention shows the water


resistance equal to or higher than that of each of the prior black dyes (comparative examples), excellent light resistance and color rendering property, especially very excellent in light resistance on ink jet papers, and the balanced water and light resistances. And, it was found that the recording fluid is excellent in storage stability and ejecting stability since the ink composition of the present invention has a high water solubility.

Effect of the Invention

[0089] The recording liquid containing the trisazo compound of the present invention is used for ink jet printing and as writing utensils. When printed on plain papers and ink jet papers, recorded images having high optical density can be formed. It has excellent light and water resistances and good storage stability as a recording liquid.



Claims

1. A trisazo compound represented by the following general formula (1):




wherein

D is an 8-hydroxy-1-naphthyl group substituted with one or two sulfo groups;
A is a group of the general formula (2) or (3):



wherein R^1 and R^2 are each independently hydrogen, amino, hydroxyl, carboxyl, sulfo, phosphono, (C1-C4)alkoxycarbonyl, (C1-C4)alkyl optionally substituted with hydroxy or (C1-C4)alkoxy, (C1-C4)alkoxy optionally substituted with hydroxy or (C1-C4)alkoxy, (C2-C4)alkanoylamino optionally substituted with hydroxy or (C1-C4)alkoxy or ureido, R^3 is hydrogen, (C1-C4)alkyl or (C1-C4)alkoxy, and R^4 is hydrogen, sulfo or phosphono;


B is a group represented by the general formula (4):



wherein R^5 , R^6 and R^7 are each independently hydrogen, hydroxyl, amino, carboxyl, (C1-C4)alkoxy, (C1-C4)alkoxycarbonyl, (C1-C4)alkyl, mono- or bis-carboxy-(C1-C3)alkylamino, carboxy-(C1-C3)alkoxy, (C1-C4)alkylamino optionally substituted with hydroxy or (C1-C4)alkoxy, (C1-C4)alkanoylamino, sulfo, halogen or ureido; and

m is 0 or 1,
or its salt.

2. A salt of a trisazo compound as claimed in claim 1 which is a lithium salt, a sodium salt, or an ammonium salt represented by the general formula (5):



wherein R^8 to R^{11} are each independently hydrogen, (C1-C4)alkyl, hydroxy-(C1-C4)alkyl or hydroxyethoxy-(C1-C4)alkyl.

3. A trisazo compound or its salt as claimed in claim 1 or 2 wherein D is

8-hydroxy-3,6-disulfo-1-naphthyl,
8-hydroxy-4,6-disulfo-1-naphthyl,
8-hydroxy-2,4-disulfo-1-naphthyl,
8-hydroxy-3,5-disulfo-1-naphthyl,
8-hydroxy-4,7-disulfo-1-naphthyl,
8-hydroxy-5,7-disulfo-1-naphthyl,
8-hydroxy-4-sulfo-1-naphthyl,
8-hydroxy-5-sulfo-1-naphthyl, or
8-hydroxy-6-sulfo-1-naphthyl.

4. A trisazo compound or its salt as claimed in any one of claims 1 to 3 wherein A is the group of the general formula (2) wherein R^1 and R^2 are each independently hydrogen, methyl, ethyl, methoxy, ethoxy, methoxyethoxy, amino, hydroxyl, carboxyl, sulfo, acetylamino, n-propionylamino or ureido, or the general formula (3) wherein R^3 is hydrogen, methyl or methoxy and R^4 is hydrogen or sulfo.
5. A trisazo compound or its salt as claimed in any one of claims 1 to 4 wherein A is the group of the general formula (2) wherein R^1 is methyl, methoxy or methoxyethoxy, and R^2 is hydrogen, methyl, methoxy, amino, hydroxyl, acetylamino or ureido.
6. A trisazo compound or its salt as claimed in any one of claims 1 to 4 wherein A is the group of the general formula (3) wherein R^3 is hydrogen or methoxy and R^4 is hydrogen or sulfo.
7. A trisazo compound or its salt as claimed in any one of claims 1 to 6 wherein B is the group of the general formula (4) wherein R^5 is hydrogen, methyl, methoxy, chlorine, carboxyl or sulfo, and R^6 and R^7 are each independently

hydroxyl, amino, carboxymethylamino, carboxymethoxy, acetylamino or ureido.

8. A trisazo compound or its salt as claimed in any one of claims 1 to 7 wherein m is zero.

5 9. An aqueous ink composition containing a trisazo compound or its salt as defined in any one of claims 1 to 8.

10. An article colored with a trisazo compound as defined in any one of claims 1 to 8 or an aqueous ink composition as defined in claim 9.

10 11. A colored article as claimed in claim 10 wherein coloring is effected by means of a printer.

15

20

25

30

35

40

45

50

55

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP98/01877

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl ⁶ C09B31/16, C09B69/02, C09D11/02 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int.Cl ⁶ C09B31/16 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS ONLINE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JP, 5-105833, A (Canon Inc.), April 27, 1993 (27. 04. 93), Claims & EP, 495520, A1 & EP, 495356, A1 & US, 5213614, A & US, 5215578, A	1-11
A	JP, 6-73320, A (Mitsubishi Kasei Corp.), March 15, 1994 (15. 03. 94), Claims (Family: none)	1-11
A	JP, 62-257971, A (CIBA-Geigy AG.), November 10, 1987 (10. 11. 87), Claims (Family: none)	1-11
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search July 13, 1998 (13. 07. 98)		Date of mailing of the international search report July 21, 1998 (21. 07. 98)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.